


GENERAL INFORMATION

KERATOCONUS



WHAT IS KERATOCONUS?

Keratoconus is a progressive, degenerative disorder characterised by thinning of the cornea – its round, dome shape starts to bulge and become cone-like instead. This process of thinning and re-shaping is termed corneal ectasia. Keratoconus is one type of corneal ectasia.

The cornea helps to bend (refract) and focus light rays onto the retina. Any abnormality in its shape leads to visual problems. These include blurred vision and difficulties seeing objects at a distance and close-up. The more advanced the keratoconus, the more severe the visual distortion.

The cause of keratoconus is not completely understood. There may be a genetic component, which is currently under investigation. Avoid continuous or vigorous rubbing of the eyes as this may be a trigger for the condition in susceptible individuals. Keratoconus has also been linked to other medical conditions, such as glaucoma, hay fever and sleep apnoea.


**KERATOCONUS GENERALLY
PROGRESSES FASTER IN YOUNGER
PATIENTS, SO EARLY DETECTION AND
TREATMENT IS VERY IMPORTANT.**





What are the symptoms of keratoconus?

- Sensitivity to light
- Seeing halos or 'ghosting'
- Distorted, blurry vision
- Difficulty driving at night
- Double vision in one eye or noticeably worse vision in one eye
- Eye strain, eye pain
- Headaches



How do you diagnose keratoconus?

Your ophthalmologist may use a number of tools to make a diagnosis of keratoconus, including:

- A **slit lamp**, which combines an intense light source with a microscope to examine your eyes
- A **keratometer** to measure the curvature of the cornea
- **Corneal topography**, which creates a 3D model of the cornea to detect any subtle changes.

HOW DO YOU TREAT KERATOCONUS?

In the early stages of keratoconus, the only treatment required may be prescription glasses to correct your vision.

Unfortunately, this is a progressive condition so your vision will eventually deteriorate, sometimes quite rapidly. As the disease progresses, other treatments may be recommended.

CONTACT LENSES

Rigid (hard) or hybrid contact lenses may be prescribed and are particularly effective in treating keratoconus for a period of time. These are made from a special material that allows the contact lens to mask the abnormal shape of the cornea and improve vision. However, contact lenses do not stop the condition from progressing and will eventually become ineffective.



CORNEAL RING SEGMENTS

This is a surgical option involving the insertion of clear plastic segments into the cornea. These segments are designed to reshape the front surface of the eye, thus correcting refractive errors caused by keratoconus.

Corneal ring segments are reserved for advanced cases of corneal ectasia, where the patient's vision is not correctable with glasses or contact lenses.




CORNEAL COLLAGEN CROSS-LINKING

Early treatment with collagen cross-linking can slow or sometimes even stop progression of keratoconus. Collagen and riboflavin (vitamin B2) are used to significantly strengthen the rigidity of the cornea.

First, the top layer of the cornea (known as the epithelium) is gently removed. The cornea is then saturated with a solution of collagen and riboflavin, which is then activated with UV light so the collagen strands bond across the cornea and strengthen it.

The procedure takes approximately 1 hour and is performed as an outpatient procedure in the clinic. Patients may experience some mild discomfort in the immediate post-operative period.

Following treatment, the patient is fitted with a contact lens that stays in place for up to 3 days.



ANTIBIOTIC DROPS ARE APPLIED TO THE TREATED EYE UNTIL ITS SURFACE HAS HEALED. THIS IS FOLLOWED BY STEROID DROPS FOR APPROXIMATELY 5 TO 6 WEEKS.

CORNEAL TRANSPLANTATION (KERATOPLASTY)

This will only be suggested if all other treatments options have been exhausted. About 10–20% of patients eventually require corneal transplantation.

There are two types of corneal transplants – partial-thickness and full-thickness (also known as penetrating). Full-thickness is usually recommended for patients with keratoconus.

A corneal transplant is a complex procedure and requires admission to a day surgery. It is generally performed under local anaesthetic, with the option of a sedative. During the procedure, your surgeon will remove the abnormal section of cornea and replace it with donor cornea, which will be stitched into place. The stitches will be removed at a later date. Your own corneal cells will gradually grow and fuse to the donor tissue. Full recovery can take up to 1 year.



FAQs

What causes keratoconus?

The cause is not fully understood, although there are indications it may be a genetic condition. Continuous or vigorous eye rubbing in susceptible individuals may lead to the development of keratoconus and should be avoided. Keratoconus has also been linked to other medical conditions, such as glaucoma, hay fever and sleep apnoea.

How common is keratoconus?

Keratoconus is usually considered a rare condition, with previous studies suggesting it affects 1 in every 2000 people. However, recent studies indicate that the incidence could be much higher – perhaps as much as 1 in 50 people. Both sexes are equally affected and most cases are diagnosed between the ages of 15 to 30 years.

Does keratoconus affect both eyes?

Generally yes, although one eye tends to be worse than the other.

Does keratoconus cause blindness?
















Severe cases may lead to legal blindness, but it's not typical for a person to become totally blind from this condition. Current treatment options allow people with keratoconus to lead relatively normal lives. Corneal transplantations are reserved for patients who no longer respond to other treatments.

Is there a Medicare rebate for corneal cross-linking?

If you have keratoconus, corneal collagen cross-linking may be recommended by your doctor. Patients undergoing this procedure are eligible for a Medicare rebate.

COMPREHENSIVE EYE CARE

Vision Eye Institute is the leading provider of ophthalmic services in Australia. Our team of highly regarded doctors includes general ophthalmologists, as well as those who specialise in specific areas/conditions of the eye.

LASER EYE SURGERY		CATARACT & LENS SURGERY	
GLAUCOMA		DIABETIC EYE DISEASE	
MACULAR DEGENERATION		RETINAL CONDITIONS	
KERATOCONUS		DRY EYE	
PTERYGIUM		CORNEAL TRANSPLANTATION	
CORNEAL CONDITIONS		NERVE-RELATED VISION PROBLEMS	
RECONSTRUCTIVE EYE SURGERY		CHILDREN'S EYE HEALTH	
GENERAL EYE HEALTH			

OUR CORNEAL CLINICS

For more information or to find a Vision Eye Institute clinic that treats keratoconus, visit:

visioneyeinstitute.com.au/services/keratoconus/

All medical and surgical procedures have potential complications. Check with your doctor before proceeding.



visioneyeinstitute.com.au



Transforming lives. That's our vision.