


enVision

22nd ANNUAL OPTOMETRY CONFERENCE



Conference roundup

Our 22nd Annual enVision Optometry Conference featured a range of clinically relevant presentations from a number of our New South Wales doctors along with our guest speaker, Dr Kate Clezy. Below we outline the key take-home messages from each presentation.



Dr Kate Clezy

Infection and optometric practice: what you don't know could hurt you

- **Means of transmission of organisms:** direct contact, droplet and aerosols
- Surfaces and clothes become contaminated with microorganisms
- **Standard precautions:** hand hygiene, personal protective equipment, sharp use and disposal, cleaning, reprocessing of reusable instruments, aseptic non-touch technique, waste management
- Perform hand hygiene before and after you touch the patient
- **Gloves:** not recommended for routine care; poor hand hygiene is common, resulting in contamination upon glove removal
- **Eye drops can be contaminated:** use single dose where appropriate
- **Disinfect equipment according to the manufacturer's instructions:** clean tonometer probes by washing/rinsing; disinfect with sodium hypochlorite; rinse; dry; store
- **Antibiotic resistance is an increasing problem:** use antibiotics only when necessary; choose narrow spectrum where possible; check treatment duration



Dr Colin Chan

The KISS guide to dry eye management in optometric practice

- Grade dry eye with targeted questionnaire on symptoms and 3 key signs: corneal staining, Schirmer's and TFBUT
- All grades of dry eye should have step 1 treatments: warm compresses, regular AFTs and omega-3 supplements
- Treatment with step 2 measures (e.g. FML, cyclosporine, IPL, tetracyclines) needs to co-exist with ongoing step 1 treatments



Dr Simon Chen

Retinal detachments: everything you need to know!

- **Types of RD:** rhegmatogenous, tractional, exudative, combined
- **Risk factors for RD:** myopia, cataract surgery, family history, RD in other eye, lattice degeneration, retinal tears, trauma, age 40–70, Stickler syndrome, Marfan syndrome
- **Evidence base:** lacking for optimal management of patients with asymptomatic retinal breaks and lattice degeneration. Management decisions regarding prophylactic laser treatment are based on individual risk/benefit analysis considering patient preference, ability to report symptoms, access to high quality urgent eye care, presence of risk factors for RD
- **Progression of retinal tears and retinal detachments**
 - Can take hours to years
 - Rate depends on symptoms, type, size, location, duration of retinal breaks, refractive error, vitreous detachment, patient age, volume of subretinal fluid present

- A young hypermetropic patient without a PVD, with a single, small, round retinal hole located inferiorly will progress slowly, if at all
- A 70-year-old high myope with multiple, large, superior horseshoe tears with a PVD and 6 disc diameters of subretinal fluid is likely to progress very quickly
- **Tips for urgent RD referrals**
 - Ensure the patient understands urgency and risk of blindness without treatment
 - Advise the patient not to eat for 6 hours or drink for 2 hours before potential surgery
- **Treatment options for RD**
 - Observation, barrier laser treatment, pneumatic retinopexy, scleral buckling surgery, vitrectomy surgery, combined vitrectomy and scleral buckling



Dr Jason Cheng

The role of MIGS in the care of glaucoma patients

- MIGS: surgical micro-devices that lower IOP with a relatively low-risk profile
- Can be inserted into 3 locations: Schlemm's canal (iStent and Hydrus), subconjunctival space (XEN) or suprachoroidal space (CyPass)
- Schlemm's canal MIGS are most effective in POAG, pseudo-exfoliative glaucoma and ocular hypertension patients who have target pressure ~18 mmHg, controlled on one medication
- Currently, iStent and Hydrus must be combined with cataract surgery to gain insurance or government rebate
- Hyphaema is a common post-operative finding that is almost always self-limiting



Dr Tess Huynh

Keratoconus: managing patients at risk


- **Keratoconus:** Bilateral, progressive, non-inflammatory
- **Genetic predisposition:** family history, environmental-atopy and eye rubbing
- **Treatment:** spectacles and contact lenses, collagen cross-linking, intrastromal rings, corneal transplant, antihistamine
- **Allergen-specific exposure:** associated with high IgE and eye rubbing, predisposes to progression of keratoconus

Collagen cross-linking: an update

- **Collagen cross-linking:** additional covalent bonding between collagen molecules has a 'stiffening' effect concentrated in anterior 200–300 µm of cornea – halts progression of keratoconus; safe
- **Intrastromal corneal rings:** remodel the cornea, regularise topography, centralise the corneal apex and reduce refraction and aberrations, improve VA and CL tolerance

enVision

22nd ANNUAL OPTOMETRY CONFERENCE



Dr Mark Jacobs

Ready, steady, go!
When to initiate treatment for POAG

- Glaucoma is an optic neuropathy
- **Types of glaucoma:** open angle, closed angle, uveitic, neovascular, congenital
- **Open angle glaucoma** suspects have at least one feature of the disease: raised IOP, suspicious optic disc appearance, repeatable visual field abnormality, strong family history
- **IOP:** most patients with raised IOP do not develop glaucoma; not all patients who develop glaucoma have high IOP
- **Physiological cupping:** bilateral; follows ISNT rule; OCT – normal RNFL and ganglion cell thickness
- **Disc area (OCT):** small <1.6 mm; medium 1.6–2.8 mm; large >2.8 mm
- **Thin CCT is a risk factor for glaucoma:** thick is protective; normal 520–580 µm
- **Family history increased risk:** parent – 2.2 x higher, sibling – 3.7 x higher
- **Treatment:** medical, SLT
- **Target IOP:** 30–50% lower than pre-treatment; consider patient lifespan, severity, family history



Dr David Ng

Diabetic retinopathy: what to look for, when to refer

- Patients should be reviewed every 1–2 years if they have no DR
- Patients with mild DR should be reviewed annually
- Patients with gestational diabetes do not need review
- Patients should be referred to an ophthalmologist if there are signs of:
 - Diabetic maculopathy and/or oedema
 - Moderate-to-severe non-proliferative DR
 - Proliferative DR
- Patients must understand that:
 - Diabetes causes blindness
 - They can be completely asymptomatic and then go blind overnight
 - They need regular review even if they have had no symptoms
 - They need to carefully control their blood sugar levels as well as their cholesterol and blood pressure, if required



A/Prof Tim Roberts

Clinical pearls in red eye management

- **Toxic epitheliopathy following cataract surgery**
 - Common cause of irritation and blurred vision 4 weeks after surgery
 - Reassure patient; use preservative free lubricants; wait for resolution before glasses prescribed
- **Bacterial endophthalmitis following cataract surgery**
 - Urgent referral: every hour counts
 - History: recent cataract surgery
 - Patient has blurred vision/pain/hypopyon/inflammation
- **Retained lens material**
 - Must be excluded in any post-cataract-surgery patient who presents with symptoms of blurred vision/raised IOP/inflammation
- **Sub-tarsal foreign body**
 - Classic vertical corneal staining
 - Red/sore/slight blur sometimes
 - Resolves upon FB removal
- **Allergic conjunctivitis**
 - Symptoms: red, itchy, watery eyes
 - Conjunctival papillae
 - Treatment with cold packs and antihistamines or mast-cell stabilisers
- **Viral follicular conjunctivitis**
 - Symptoms: red, watery eyes
 - Follicles not papillae, clear non-mucopurulent discharge
 - Treat with supportive lubricants, not topical antibiotic
- **Acute uveitis**
 - Symptoms: bilateral redness, tearing, sensitivity to light, painful
 - AC activity and some disc swelling
 - Urgent referral, treated with topical steroids and cyclopegics as well as systemic management
- **HSV1 keratitis**
 - Slight blurring of vision/redness/sore eye
 - Typical dendritic corneal lesions
- **Marginal keratitis**
 - Red/painful eye/light-sensitive and some blur and watery discharge
 - History: blepharitis with inflamed lid margins
 - Hypersensitive reaction to staphylococcus endotoxins
 - Be suspicious of bacterial keratitis
 - Treat with lid toilette, antibiotic and steroid drops



Prof Gerard Sutton

SMILE – an update

- An essential option for patients considering myopic refractive surgery
- My preferred option for higher levels of myopia >3 D
- Causes less dry eye than LASIK or ASLA/PRK
- LASIK is still preferred for lower levels of myopia and moderate astigmatism
- ASLA/PRK is preferred for thinner corneas or questionable topography

Fuchs' endothelial dystrophy – a guide for optometrists

- A common problem
- New types of endothelial keratoplasty such as DMEK are providing improved visual outcomes
- Femtosecond laser reduces endothelial cell loss in cataract surgery
- Cataract surgery in patients with corneal issues (e.g. Fuchs' dystrophy, keratoconus) requires a nuanced approach to treatment to achieve optimal outcomes



Dr Patrick Versace

Demystifying post-cataract surgery symptoms & signs

- **Post-op cataract surgery:** review day 1, 1 week, 3 months, 1 year, 10 years
- **AC activity:** always at day 1, common at 1 week, rare at 1 month; rebound at 6 weeks as finish post-op drops, treat with steroids and non-steroidals (Ilevro). Persistent iritis post-cataract can be infectious or inflammatory
- **Steroids can cause cataract or rise in IOP:** less IOP rise with FML
- Can get post-cataract corneal oedema or Descemet's folds
- **Warning signs after cataract surgery:** vision loss, pain, redness
- **Endophthalmitis:** pain, redness, loss of vision, inflammation, hypopyon, cells in vitreous
- **Consider IOL:** position, refractive outcome, toric axis, PC, clarity
- **PCO:** treated with YAG laser
- **Dysphotopsia caused by lens edge:** positive or negative

eyeMatters

visioneyeinstitute.com.au

vision
eye institute